

DC49257 | DLin-K-C3-DMA

Instructions for Use

DLin-KC3-DMA, a nucleic acid, shows in vivo silencing activity. DLin-K-C3-DMA can be used in the synthesis of nucleic acid-lipid particle to delivery of nucleic acid.

Product information

CAS No.	Not available
Chemical Name	DLin-K-C3-DMA
Formula	C44H81NO2
M.Wt	656.1195
Purity	>95%
Storage	Original product: up to 1 year under the storage conditions stated on the vial, kept tightly sealed. Stock solutions: aliquots at -20°C, generally up to one month.
Shipping condition	Ships from Shanghai. Pure lipid is stable during ice-pack transport.

Formulation & study context

Literature values apply to the cited study only. Administration routes describe research, not clinical instructions.

A complete product-specific formulation has not been verified. No standard ratio is substituted.

N/P ratio	Not reported in the verified record
Cargo	Not available in the verified record
Administration route	Not available in the verified record
Animal model	Not available in the verified record
Formulation process	Not available in the verified record

N/P is a nitrogen-to-phosphate molar ratio. Lipid:cargo mass ratio is a different quantity and must not be used as N/P.

Stock-solution preparation table

Calculated using product-page MW: 656.1195 g/mol. Volumes are final solution volumes.

Mass	1 mM	5 mM	10 mM
10 mg	15.241 mL	3.048 mL	1.524 mL
25 mg	38.103 mL	7.621 mL	3.810 mL
50 mg	76.206 mL	15.241 mL	7.621 mL

Mass	1 mM	5 mM	10 mM
100 mg	152.411 mL	30.482 mL	15.241 mL
250 mg	381.028 mL	76.206 mL	38.103 mL

Volume (mL) = mass (mg) × 1000 / [MW (g/mol) × concentration (mM)]. Mathematical conversions only; not measured solubility. Confirm solvent compatibility and the exact material before preparation.

Handling & storage

Provided that storage conditions are as stated on the product vial and the vial is kept tightly sealed, the product can be stored for up to 1 year.

Wherever possible, prepare and use solutions on the same day. If stock solutions must be prepared in advance, store them as aliquots in tightly sealed vials at -20°C. Generally, these will be usable for up to one month.

Before use, and prior to opening the vial, allow the product to equilibrate to room temperature for at least 1 hour.

References

Not available