

DC60483 | LIS10W

Instructions for Use

LIS10W is a sugar-alcohol-derived ionizable lipid with L-sorbitol as the precursor.

Product information

CAS No.	Not available
Chemical Name	LIS10W
Formula	C72H144O12N2
M.Wt	1,229.95
Purity	>98%
Storage	Original product: up to 1 year under the storage conditions stated on the vial, kept tightly sealed. Stock solutions: aliquots at -20°C, generally up to one month.
Shipping condition	Ships from Shanghai. Pure lipid is stable during ice-pack transport.

Formulation & study context

Literature values apply to the cited study only. Administration routes describe research, not clinical instructions.

A complete product-specific formulation has not been verified. No standard ratio is substituted.

N/P ratio	Not reported in the verified record
Cargo	Not available in the verified record
Administration route	Not available in the verified record
Animal model	Not available in the verified record
Formulation process	Not available in the verified record

N/P is a nitrogen-to-phosphate molar ratio. Lipid:cargo mass ratio is a different quantity and must not be used as N/P.

Stock-solution preparation table

Calculated using product-page MW: 1229.95 g/mol. Volumes are final solution volumes.

Mass	1 mM	5 mM	10 mM
10 mg	8.130 mL	1.626 mL	0.813 mL
25 mg	20.326 mL	4.065 mL	2.033 mL
50 mg	40.652 mL	8.130 mL	4.065 mL
100 mg	81.304 mL	16.261 mL	8.130 mL

Mass	1 mM	5 mM	10 mM
250 mg	203.260 mL	40.652 mL	20.326 mL

Volume (mL) = mass (mg) × 1000 / [MW (g/mol) × concentration (mM)]. Mathematical conversions only; not measured solubility. Confirm solvent compatibility and the exact material before preparation.

Handling & storage

Provided that storage conditions are as stated on the product vial and the vial is kept tightly sealed, the product can be stored for up to 1 year.

Wherever possible, prepare and use solutions on the same day. If stock solutions must be prepared in advance, store them as aliquots in tightly sealed vials at -20°C. Generally, these will be usable for up to one month.

Before use, and prior to opening the vial, allow the product to equilibrate to room temperature for at least 1 hour.

References

Not available