

DC80071 | A18-Iso5-2DC18

Instructions for Use

A18-Iso5-2DC18 that could not only deliver mRNA vaccines robustly but also activate the stimulator of interferon genes (STING) pathway.

Product information

CAS No.	Not available
Chemical Name	A18-Iso5-2DC18
Formula	C50H93N3O2
M.Wt	768.29
Purity	>98%
Storage	Original product: up to 1 year under the storage conditions stated on the vial, kept tightly sealed. Stock solutions: aliquots at -20°C, generally up to one month.
Shipping condition	Ships from Shanghai. Pure lipid is stable during ice-pack transport.

Formulation & study context

Literature values apply to the cited study only. Administration routes describe research, not clinical instructions.

A complete product-specific formulation has not been verified. No standard ratio is substituted.

N/P ratio	Not reported in the verified record
Cargo	Not available in the verified record
Administration route	Not available in the verified record
Animal model	Not available in the verified record
Formulation process	Not available in the verified record

N/P is a nitrogen-to-phosphate molar ratio. Lipid:cargo mass ratio is a different quantity and must not be used as N/P.

Stock-solution preparation table

Calculated using product-page MW: 768.29 g/mol. Volumes are final solution volumes.

Mass	1 mM	5 mM	10 mM
10 mg	13.016 mL	2.603 mL	1.302 mL
25 mg	32.540 mL	6.508 mL	3.254 mL
50 mg	65.080 mL	13.016 mL	6.508 mL

Mass	1 mM	5 mM	10 mM
100 mg	130.159 mL	26.032 mL	13.016 mL
250 mg	325.398 mL	65.080 mL	32.540 mL

Volume (mL) = mass (mg) × 1000 / [MW (g/mol) × concentration (mM)]. Mathematical conversions only; not measured solubility. Confirm solvent compatibility and the exact material before preparation.

Handling & storage

Provided that storage conditions are as stated on the product vial and the vial is kept tightly sealed, the product can be stored for up to 1 year.

Wherever possible, prepare and use solutions on the same day. If stock solutions must be prepared in advance, store them as aliquots in tightly sealed vials at -20°C. Generally, these will be usable for up to one month.

Before use, and prior to opening the vial, allow the product to equilibrate to room temperature for at least 1 hour.

References

Not available